




Patient and procedure risk stratification backed by clinical evidence

While surgical patients may benefit from Prevena Therapy, patients at high risk for complications such as surgical site infection may see added benefit. The following uses select study data¹⁻³ to provide an illustrative guide to aid in risk stratification. This is not an all-inclusive list of risk factors. Clinicians are advised to use their clinical judgment to identify high-risk patients or high-risk procedures.

Start here



References

1. Willy C, Agarwal A, Andersen CA, De Santis G, Gabriel A, Grauhan O, Guerra OM, Lipsky BA, Malas MB, Mathiesen LL, Singh DP, Reddy VS. Closed incision negative pressure therapy: international multidisciplinary consensus recommendations. *Int Wound J.* 2017 Apr;14(2):385-398. [OPEN ACCESS](#)
2. Grauhan O, Navasardyan A, Hofmann M, Muller P, Stein J, Hetzer R. Prevention of poststernotomy wound infections in obese patients by negative pressure wound therapy. *Journal of Thoracic and Cardiovascular Surgery.* 2013 May;145(5):1387-92. [OPEN ACCESS](#)
3. Suelo-Calanao RL, Thomson R, Read M, Matheson E, Isaac E, Chaudhry M, Loubani M. The impact of closed incision negative pressure therapy on prevention of median sternotomy infection for high risk cases: a single centre retrospective study. *Journal of Cardiothoracic Surgery.* 2020 Aug 19;15(1):222. [OPEN ACCESS](#)

Advancing the standard of care.



View the cardiothoracic clinical evidence summary



3M Company
2510 Conway Ave.
St. Paul, MN 55144 USA

Phone 1-800-275-4524 (NPWT products)
1-800-228-3957

Web 3m.com/medical

3M™ Prevena™ 125 Therapy Unit and 3M™ Prevena™ Plus 125 Therapy Unit manage the environment of closed surgical incisions and remove fluid away from the surgical incision via the application of -125mmHg continuous negative pressure. When used with legally marketed compatible dressings, Prevena 125 and Prevena Plus 125 Therapy Units are intended to aid in reducing the incidence of seroma; and, in patients at high risk for post-operative infections, aid in reducing the incidence of superficial surgical site infection in Class I and Class II wounds.

The effectiveness of Prevena Therapy in reducing the incidence of SSIs and seroma in all surgical procedures and populations has not been demonstrated. See full indications for use and limitations at mykci.com.

The indication statement does not apply to the Prevena Plus 125 Therapy Unit (14-Day) that comes with the 3M™ Prevena Restor™ System Kits (see Prevena Restor System Instructions for Use).

NOTE: Specific indications, limitations, contraindications, warnings, precautions and safety information exist for these products and therapies. Please consult a clinician and product instructions for use prior to application. Rx only.

© 2023 3M. All rights reserved. 3M and the other marks shown are marks and/or registered trademarks. Unauthorized use prohibited. US_70-2013-1483-1