Solventum



Your first step is wound bed prep

Let's make every CTP application count

Start with wound bed prep for healing success

With limited cellular and tissue based product (CTP)/skin sub applications, preparation is more critical than ever

Hard-to-heal wounds, often stuck in a prolonged inflammatory phase, fail to progress through the normal stages of healing. This results in excess matrix metalloproteinases (MMPs). which degrade the extracellular matrix and growth factors essential for tissue repair, further impairing healing.^{1,2}



Clinical guidelines recommend advanced therapies, including skin substitutes, for wounds that fail to respond to initial treatments within four

When done properly, wound bed preparation (including debridement):

- · Reduces bacterial load
- · Removes necrotic tissue
- Lowers inflammatory proteases

All of these are critical steps to promote healing and facilitate the successful application of advanced therapies. 1,6,7

"Appropriate wound bed preparation is vital for optimal patient outcomes with skin substitutes."1

Mulder et al, 2024



Did you know?

In a recent expert panel consensus, 100% of panel members concluded that the combination of 3M™ Promogran Prisma™ Collagen Matrix with ORC and Silver and 3M[™] ActiV.A.C.[®] Therapy System with 3M[™] V.A.C.[®] Granufoam Dressing may help:

- Remove barriers to healing
- Balance the wound environment
- Promote granulation tissue development⁸

By integrating advanced therapies like Promogran Prisma Matrix and ActiV.A.C. Therapy System, clinicians can address complex wound types, including diabetic ulcers, venous leg ulcers, pressure injuries, surgical wounds and more, showing benefits such as:

- Faster time to granulation tissue development
- Reduced wound area and volume
- Improved wound bed preparation outcomes^{8,9}

A consensus panel recommendation outlined the use of Promogran Prisma Matrix with ActiV.A.C. Therapy System and V.A.C.® Granufoam Dressings to be considered at the earliest opportunity once deemed clinically appropriate.8

CASE STUDY

DFU treated with 3M™ Promogran Prisma™ Collagen Matrix with ORC and Silver and 3M[™] ActiV.A.C.[™] Therapy System³







Images used with permission from Robert Klein, DPM.

Silver collagen oxidized regenerated cellulose (ORC) dressing with negative pressure wound therapy (NPWT) and reticulated open cell foam (ROCF) dressing use in an infected diabetic neuropathic ulcer.

- A. Diabetic neuropathic ulcer of the heel at presentation.
- B. The wound after abscess drainage, sharp debridement, and initiation of antibiotic therapy.
- C. The wound after 28 days of combination therapy use.

A powerful pair for wound bed prep



Clinical evidence for 3M[™] Promogran Prisma™ Collagen Matrix with ORC and Silver:

- Increased wound closure rates: A meta-analysis showed that wounds treated with collagen/ORC dressings were 1.74 times more likely to close compared to standard dressings (n=1521).10
- Fewer withdrawals due to infection: An RCT showed fewer withdrawals because of infection with Promogran Prisma Matrix.¹¹

Promogran Prisma Matrix

Helps manage inflammation and deal with the inflammatory environment.¹²



Collagen - 55%

- Acts as a **sacrificial substrate** for excessive MMP¹³
- Bioresorbable¹⁴ into amino acids which can be reused by cells to make new proteins
- Encourages the movement of neutrophils, macrophages and fibroblasts, stimulating new tissue growth¹⁵

Oxidized Regenerated Cellulose (ORC) — 44%

In vitro studies have demonstrated the potential benefits of ORC12:

- Degrades to glucose and glucuronic acid, lowering the pH which helps control bacteria
- Protects growth factors from degradation
- Reduces protease activity levels, specifically elastase

Silver (Ag) — 1% Ag/ORC

Due to the antibacterial properties of silver, Promogran Prisma Matrix is known to provide an effective antibacterial barrier against common wound pathogens in vitro. 16

protect growth factors cleaning Ower and cleaning Owe prepare for closure

lower MMPs reduce inflammation microstrain a set sele **P** action promote granulation reduce edema

remove exudate

optimize healing promote perfusion macros draw wound ⇉ <u>a</u> edges together



3M[™] ActiV.A.C.[™] Therapy System with 3M™ V.A.C.® Granufoam™ Dressing

Creates macrostrain and microstrain, removes exudate and infectious material, reduces edema and promotes granulation tissue formation.

In a retrospective data analysis of 485 cases, wounds treated with ActiV.A.C. Therapy in combination with Promogran Prisma Matrix showed a 40% reduction in wound area compared to a 9% reduction with NPWT alone.9

"Wounds managed with NPWT + OCSO [oxidized collagen/ silver-ORC dressings] were significantly more likely to improve and/or heal compared with wounds that received NPWT alone."9

Hou & Griffin, 2024

3M[™] Promogran Prisma[™] Collagen Matrix with ORC and Silver and 3M[™] ActiV.A.C.[™] Therapy System

The most studied NPWT product in the world is now labeled for use with the most studied collagen dressing in the world

Combined, these two leading wound treatment therapies have been shown to support improved healing outcomes.⁹

To order ActiV.A.C. Therapy, visit 3M.com/Express



ORDERING INFORMATION

3M™ Promogran Prisma™ Collagen Matrix with ORC and Silver

Item Code	Eaches/Carton/Case	Size	HCPCS Code
MA028**	10 ea/ct - 4 ct/cs	4.34 sq. in. Hexagon	A6021
MA123**	10 ea/ct - 4 ct/cs	19.1 sq. in. Hexagon	A6022
MA032 ⁺	6 ea/ct - 4 ct/cs	3/8" x 3/8" x 12 5/8" Rope	A6024

3M™ Promogran™ Collagen Matrix with ORC

Item Code	Eaches/Carton/Case	Size	HCPCS Code
PG004 [†]	10 ea/ct - 4 ct/cs	4.34 sq. in. Hexagon	A6021
PG019 [†]	10 ea/ct - 4 ct/cs	19.1 sq. in. Hexagon	A6022



3M[™] V.A.C.[®] Peel and Place Dressing may be used to bolster skin substitutes, flaps and grafts.*

*Promogran Prisma Collagen Matrix is not labeled for use under V.A.C.® Peel and Place Dressing

**Only MA028 and MA123 are indicated for use with ActiV.A.C. Therapy System

*Not labeled for use with negative pressure

NOTE: Specific indications, contraindications, warnings, precautions and safety information exist for these products and therapies. Please consult a clinician and product instructions for use prior to application. Rx only.

References

- 1. Mulder G, Lavery LA, Marston WA et al (2024) Skin substitutes for the management of hard-to-heal wounds. London: Wounds International. Available to download from: www.woundsinternational.com
- 2. Przekora A (2020) A concise review on tissue engineered artificial skin grafts for chronic wound treatment: can we reconstruct functional skin tissue in vitro? Cells 9(7): 1622.
- 3. IWGDF (2023) Guidelines on interventions to enhance healing of foot ulcers in people with diabetes. Available at: www.iwgdfguidelines.org (accessed 31.07.2024)
- 4. Vecin NM, Kirsner RS (2023) Skin substitutes as treatment for chronic wounds: current and future directions. In: Frontiers in Medicine (Vol. 10). Frontiers Media SA.
- 5. Tettelbach W, Forsyth A (2024) Current practices using cellular, acelluar and matrix-like products (CAMPs). Br J Nurs 33(4): S4-8.
- 6. Schultz GS, Sibbald RG, Falanga V et al (2003) Wound bed preparation: a systematic approach to wound management. Wound Repair Regen 11: S1-28.
- 7. Demidova-Rice TN, Hamblin MR, Herman IM (2012) Acute and impaired wound healing: Pathophysiology and current methods for drug delivery, part 1: Normal and chronic wounds: Biology, causes, and approaches to care. Adv Skin Wound Care 25(7): 304-14.
- 8. Klein R, Desvigne MN, Greenstein E, et al. Use of silver collagen oxidized regenerated cellulose dressings in conjunction with negative pressure wound therapy: expert panel consensus recommendations. Wounds. 2024;36(8):263-273. doi:10.25270/wnds/2402
- 9. Hou Y, Griffin L. Comparative effectiveness of negative pressure wound therapy with and without oxidized regenerated cellulose (ORC) collagen/silver-ORC dressing. Poster presented at: SAWC, National Harbor, MD, 2023.
- 10. Chen Y, Du P, Lv G. A meta-analysis examined the effect of oxidized regenerated cellulose/collagen dressing on the management of chronic skin wounds. *Int Wound J.* 2023;20(5):1544-1551. doi:10.1111/iwj.14009.
- 11. Gottrup F, Cullen B, Karlsmark T, Bischoff-Mikkelsen M, Nisbet L, Gibson M. Randomized controlled trial on collagen/oxidized regenerated cellulose /silver treatment. Wound Repair & Regeneration 2013; 21: 1-10.
- 12. Cullen B, Watt PW, Lundqvist C, et al. The role of oxidized regenerated cellulose/collagen in chronic wound repair and its potential mechanism of action. Int J Biochem Cell Biol 2002; 34(12): 1544-56.
- 13. Schultz GS, Ladwig G, Wysocki A. Extracellular Matrix: a review of its roles in acute and chronic wounds. World Wide Wounds. 2005; Accessed August 19, 2020. http://www.worldwidewounds.com/2005/august/Schultz/Extrace-Matric-Acute-Chronic-Wounds.html.
- 14. Wagermaier W, Fratzl P. Collagen. In: Moeller M, Matyjaszewski K, eds. Polymer Science: A Comprehensive Reference. 1st ed. *Elsevier Focal Press Science and Tech.* 2012:35–55.
- Postlewaite AE, Seyer JM, Kang AH. Chemotactic attraction of human fibroblasts to type I, II and III collagens and collagen-derived peptides. Proc Nat Acad Sci USA. 1978;75(2):871–875.
- 16. Bourdillon KA, Delury C, Cullen B. Biofilms and delayed healing an in vitro evaluation of silver and iodine containing dressings and their effect on bacterial and human cells. International Wound Journal 2017. ISSN 1742-4801.



Solventum Corporation 2510 Conway Avenue East Maplewood, MN 55144 Phone 1-800-243-6894 1-203-237-5541 Web www.Solventum.com

Solventum, formerly 3M Health Care

© Solventum 2025. Solventum, the S logo and other trademarks are trademarks of Solventum or its affiliates. 3M, the 3M logo and other 3M owned and/or licensed trademarks are trademarks of 3M. Other trademarks are the property of their respective owners. US_70-2013-1786-7